



CAT 360

Catastrophe Risk from Every Perspective

The CAT 360 is a quarterly newsletter that features articles developed by our Research and Development Team and covers topics that relate to Catastrophe Modeling, Natural Perils and Information Technology on a global basis. Please feel free to contact the editors if you have any questions or comments regarding any of our publications.

Eds.

Second Quarter 2008

ACE Tempest Re
ACE Building
17 Woodbourne Avenue
Hamilton HM08
Bermuda

441 292-2603 tel
441 295-2888 fax
acetempetre@acegroup.com
www.acetempetre.com

Feature Stories

[Click the Article Title to Read Full Story](#)

The 2008 Atlantic Hurricane Season Forecasts **Forecast Highlights** **Discussion** **Summary**

The 2008 Atlantic Hurricane Season Forecasts A Credible Crystal Ball?

Ever since Hurricane Andrew devastated Florida in 1992, the annual Atlantic Hurricane season forecast has always stirred interest in the coastal communities in the United States. But how accurate are these forecasts and what are the factors that influence them?

This article will answer those questions with a behind the scenes discussion of the organizations that develop these forecasts along with an update to the 2008 Atlantic Hurricane Season forecast.

[^ TOP](#)

Forecast Highlights

- The three major institutes: The National Oceanic and Atmospheric Administration (NOAA), Tropical Storm Risk (TSR), and Colorado State University (CSU) have released their long-term forecast for the upcoming hurricane season. All three forecasts suggest that 2008 will be slightly to moderately above the long term average and close to the mid-term average since 1995.
- Current conditions suggest a continued weakening of La Niña through June, and a possible transition to an ENSO neutral state in June-July 2008. This is consistent with dynamic and statistic model forecasts. La Niña years are associated with above normal hurricane activity.
- According to the UK Met Office, sea surface temperatures (SSTs) in the Tropical Atlantic will be close to average in the summer. This is consistent with NOAA's Coupled Forecast System (CFS) model which predicts close to average or slightly above average SSTs in the peak hurricane months.

[^ TOP](#)

Discussion

Table 1 summarizes the forecast from the previously mentioned three institutes. Consistency of these forecasts does not guarantee an accurate forecast will occur, as evidenced in the 2006 and 2007 hurricane seasons. Model forecasting skill needs to be considered when

May, the UK Met Office's forecast will not be available until mid-June.

By comparison, the NOAA CFS model is used to develop mid to long term forecasts. This model has been shown to have reasonable forecast accuracy with

	Climatology Average	NOAA	CSU	TSR
Named Storms	10	12 -16	15	14.8
Hurricanes	6	6 - 9	8	7.8
Major Hurricanes	2.7	2 - 5	4	3.5


Table 1 - Forecast for the 2008 Atlantic Hurricane Season

interpreting any forecast. A forecast with marginal accuracy may be no better than using the average or using a pure guess.

The forecast scheme at TSR has not changed from prior years. It continues to use forecasted trade winds and sea surface temperatures. The trade wind attribute is highly correlated with the current ENSO condition and is therefore reflected in their forecast. CSU changed its forecast scheme for the 2007 season, and has once again changed it for 2008. Its methodology now uses data from 1950 to 2007 in developing a linear regression. Because it is newly implemented, a testing period is not available to verify this new method.

Last year, the UK Met office drew much attention when it predicted a normal to below normal hurricane season. This was in contrast to the other institutes which predicted an above normal hurricane season for 2007. One major difference is that the UK Met office uses a dynamic ocean model to forecast the SSTs in the summer months. By June of 2007, the dynamic model suggested below average SSTs for the summer months, which was a driving factor for its below normal forecast. This turned out to be the most accurate forecast of the actual season.

UK Met Office Summer SST Outlook (Source: UK Met Office) 2008 in Comparison with the Summer of 2007




April 2008


a three-month lead time for certain parameters. The three-month average SST anomalies are in the range of 0 to 0.5°C. On average, the SST anomalies are about 0.1°C. However, hurricane genesis and intensification are not solely decided by the SST. Low vertical shear accompanying a La Niña event creates a favorable environment for intensification. The correlation skill of the CFS model forecast in the Tropical Atlantic using this lead time is about 0.3 - 0.4. The forecast is more reliable when using a lead time of 1 - 2 months and can increase to 0.8 in the tropics. Interestingly, the CFS model also predicts the dissipation of the current La Niña condition by June and even a shift to above normal SSTs in the Nino 3.4 region (the Pacific off the West South American Coast) in the summer. In contrast, the UK Met Office predicts a longer La Niña event.

The ENSO predictability model is not as accurate in the early part of the year compared to the summer months since the ENSO indicators tend to change sign in the spring. This is commonly known as the spring barrier. Nevertheless, the spread of model results and the similar historical years suggests that a neutral ENSO is very likely this summer. The correlation accuracy with a three month lead forecast is above 0.85.

The Bermuda High is also an important factor in deciding the track of hurricanes. Last year a southerly positioned Bermuda High directed the storms in the Caribbean and Mexico. As a result, the US mainland escaped major hurricane landfalls. In contrast, the position of the 2004 Bermuda High caused three hurricanes (Charley, Frances and Jeanne) to hit a similar region in Florida. The position of the Bermuda High is correlated with the strength of the Northern Atlantic Oscillation (NAO). Negative NAO conditions tend to lead to more US landfalling storms. Unfortunately, forecasting the summer NAO is challenging, with a forecast beyond two weeks considered to be unreliable.




May 2007



June 2007

The figures above are the UK Met Office's SST forecasts at different times. As of April 2008, the SST in the Tropical Atlantic is expected to be close to average. You can see that compared to May 2007, June 2007 was cooler (areas of green in the Caribbean). Generally speaking, forecasts with a shorter lead time are more credible and in this case, it did make a significant difference. Unlike the other three institutes which release their forecasts in



Observed weekly SST anomalies with respect to 1971-2000. (Source: NOAA CPC)



Current ENSO condition in comparison with similar years in the past. (Source: IRI) Index value persistently less than -0.5 indicates La Niña, while index value persistently greater than +0.5 indicates El Niño.

[^ TOP](#)

Summary

While science has made great advancements in understanding the conditions that effect the formation of hurricanes, forecasting the upcoming hurricane season is still as much art as it is science. Information provided by these organizations are state of the art and highly valuable in preparing the public for severe hurricane events.

Whether the hurricane forecasts are accurate or completely wrong, the scientific community will continue to gain invaluable knowledge with the completion of every hurricane season and apply what they learn for years down the road.

[^ TOP](#)

The information contained in these newsletters was compiled by ACE Tempest Re and is intended as background information only. All information is provided "as is" with no guarantees of completeness, accuracy or timeliness and without warranties of any kind, express or implied. ACE Tempest Re is not responsible for, and expressly disclaims all liability for, damages of any kind, whether direct or Indirect, consequential, compensatory, actual, punitive, special, incidental or exemplary, arising out of use, reference to, or reliance on any information contained herein.

This is a message from ACE Tempest Re, ACE Building, 17 Woodbourne Avenue, Hamilton HM08, Bermuda.

© 2009 ACE Tempest Re | [Terms of Use](#) | [Company Licensing](#)

This email was sent to: shoopn@digitlev.com

This email was sent by: ACE Tempest Re
ACE Building, 17 Woodbourne Avenue, Hamilton HM08 Bermuda

We respect your right to privacy - [view our policy](#)